

# Modelling the near-Earth magnetospheric field asymmetries

Martin Fillion<sup>1,2</sup>, Arnaud Chulliat<sup>1,2</sup>, Patrick Alken<sup>1,2</sup>, Mikhail Kruglyakov<sup>3</sup>, Alexey Kuvshinov<sup>4</sup>

<sup>1</sup>CIRES, University of Colorado Boulder <sup>2</sup>NOAA National Centers for Environmental Information, <sup>3</sup>Department of Physics, University of Otago, New Zealand <sup>4</sup>Institute of Geophysics, ETH Zürich, Switzerland

## Introduction

Data-based modeling of the near-Earth magnetospheric field is challenging due to the highly dynamical nature of magnetospheric currents and to the sparse available data. One important characteristic of this field is the so-called local time asymmetry, which can be clearly observed during the main phase of geomagnetic storms. Modeling this asymmetry requires a dataset with relatively good spatial coverage at a relatively high time resolution. The network of ground magnetic observatories provides the only magnetic data that meet these criteria. In this study, we present a data-based model of the near-Earth magnetic field produced by electric currents in the inner magnetosphere.




Figure 1: sketch of magnetospheric currents

## Data preprocessing

We use ground observatory hourly magnetic field data from the World Data Center database between 5° and 55° of quasi dipole latitude from 1997 to 2021. The data are corrected from the following fields:

- Main field using the CHAOS 7.13 model<sup>3</sup>
- Crustal biases from Califf et al. (2022)
- E-region mid- and low-latitude ionospheric field using the DIFI model<sup>2</sup>
- Static GSM magnetospheric field using the CHAOS 7.13 model<sup>3</sup>





Figure 2: Position of geomagnetic observatories providing data used in this study.

**Methodology:** We derive a global spherical harmonics model of the external field every one-hour timestamp:

- Select all available data for one timestamp
- Solve an inverse problem to find the best spherical harmonics model in dipole magnetic coordinates
- The spherical harmonics coefficients are separated in their external and induced parts using an approach developed by Kruglyakov et al. (2022)
- Repeat process for next timestamp



## Methodology and results




Figure 3: Comparison between results obtained in a study of Le et al. (2011) using data from the C/NOFS satellite (6 panels on the left) and results obtained using our spherical harmonics model (6 panels on the right) for the geomagnetic storm of July 2009. Each panel corresponds to one timestamp. The angle shows the local time and the radial distance from the dashed black circle (left) or the black plain circle (right) the magnitude of the magnetic signal in nT. The Dst index – constant through all local times – is shown with a blue circle (left) or an orange circle (right).

Figure 4: (Upper) Root Mean Square Error (RMSE) between the time series of the South component of the Swarm satellite vector data and the corresponding time series of the South component of the model prediction as a function of the Hp30 index<sup>6</sup> for a degree 2 model (blue), a degree 3 model (purple), a modified CHAOS 7.13 model with an in-house implementation of the RC index (red), and the official CHAOS 7.13 index (green). (Bottom) Relative difference between the RMSE of the modified CHAOS 7.13 model and the RMSE of the degree 2 model (blue), of the degree 3 model (purple) and of the official CHAOS 7.13 (green).

$$\text{Reldiff} = \frac{\text{RMSE}_{\text{RC}} - \text{RMSE}}{\text{RMSE}_{\text{RC}}} \times 100$$

**Observation:** Degree 2 and 3 models improve fit to Swarm data in comparison to the CHAOS 7.13 model on the South component above Hp30=2.




Figure 5: Average South component - in dipole magnetic coordinate - over 24 years predicted by the non-zonal terms of the spherical harmonics expansions - corresponding to the asymmetric part of the magnetospheric field - at the magnetic equator, and as a function of the Kp index and local time.

**Observation:** Figure 5 shows a relatively strong dawn-dusk asymmetry for Kp above 3 and a weaker noon-midnight asymmetry for Kp below 2.

## Conclusions

To first order, the magnetospheric local time asymmetry can be represented by a degree 2 spherical harmonics external potential field model. This approach can be used for at least two distinct purposes:

- Provide a better correction of the magnetospheric field for core field modelling
- Provide constraints for the study of geomagnetic storms and substorms

In the future, the model will be improved by developing an algorithm to minimize contamination from the ionospheric field and to balance potential artefacts associated with the heterogeneous geographic data coverage.

<sup>1</sup>Califf, Samuel, Patrick Alken, Arnaud Chulliat, Brian Anderson, Kenneth Rock, Sarah Vines, Robin Barnes, and Kan Liou. "Investigation of Geomagnetic Reference Models Based on the Iridium constellation." *Earth, Planets and Space* 74, no. 1 (February 25, 2022): 37. <https://doi.org/10.1186/s40623-022-01574-w>.

<sup>2</sup>Chulliat, A., P. Vigneron, and G. Hulot. "First Results from the Swarm Dedicated Ionospheric Field Inversion Chain." *Earth, Planets and Space* 68, no. 1 (December 2016): 104. <https://doi.org/10.1186/s40623-016-0481-6>.

<sup>3</sup>Finlay, Christopher C., Clemens Kloss, Nils Olsen, Magnus D. Hammer, Lars Tøffner-Clausen, Alexander Grayver, and Alexey Kuvshinov. "The CHAOS-7 Geomagnetic Field Model and Observed Changes in the South Atlantic Anomaly." *Earth, Planets and Space* 72, no. 1 (December 2020): 156. <https://doi.org/10.1186/s40623-020-01252-9>.

<sup>4</sup>Kruglyakov, Mikhail, Alexey Kuvshinov, and Manoj C. Nair. "A Proper Use of the Adjacent Land-Based Observatory Magnetic Field Data to Account for the Geomagnetic Disturbances during Offshore Directional Drilling." *Earth and Space Science Open Archive*, October 13, 2022. World. <https://doi.org/10.1002/essoar.10512088.2>.

<sup>5</sup>Le, Guan, William J. Burke, Robert F. Pfaff, Henry Freudenreich, Stefan Maus, and Hermann Lühr. "C/NOFS Measurements of Magnetic Perturbations in the Low-Latitude Ionosphere during Magnetic Storms." *Journal of Geophysical Research: Space Physics* 116, no. A12 (2011). <https://doi.org/10.1029/2011JA017026>.

<sup>6</sup>Yamazaki, Y., Matzka, J., Stolle, C., Kervalishvili, G., Rauberg, J., Bronkalla, O., Morschhauser, A., Bruinsma, S., Shprits, Y. Y., & Jackson, D. R. (2022). Geomagnetic Activity Index Hpo. *Geophysical Research Letters*, 49(10), e2022GL098860. <https://doi.org/10.1029/2022GL098860>